

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Ambient Air Pollution, Placental Mitochondrial DNA Content, and Birth Weight in the INMA (Spain) and ENVIRONAGE (Belgium) Birth Cohorts

Diana B.P. Clemente, Maribel Casas, Nadia Vilahur, Haizea Begiristain, Mariona Bustamante, Anne-Elie Carsin, Mariana F. Fernández, Frans Fierens, Wilfried Gyselaers, Carmen Iñiguez, Bram G. Janssen, Wouter Lefebvre, Sabrina Llop, Nicolás Olea, Marie Pedersen, Nicky Pieters, Loreto Santa Marina, Ana Souto, Adonina Tardón, Charlotte Vanpoucke, Martine Vrijheid, Jordi Sunyer, and Tim S. Nawrot

Table of contents

Table S1. Data from each campaign and predictor variables of air pollution in each cohort

Figure S1. GAM models that show the relation between (A) NO₂ exposure (µg/m³) during the entire pregnancy and mtDNA content, (B) NO₂ exposure (µg/m³) during the entire pregnancy and birth weight (g), and (C) mtDNA content and birth weight (g). This model provides p-values for the null hypothesis of no linearity. 1: ENVIRONAGE, 2: INMA

Table S2. Percent change in placental mtDNA content in association with prenatal NO₂ exposure in the four different INMA subcohorts

Table S3. Association between pregnancy average NO₂ exposure and birth weight in the four different INMA subcohorts

Table S4. Mediation analysis of the estimated effects of prenatal NO₂ exposure (µg/m³) on birth weight through placental mtDNA content in the pooled sample and the boys of the pooled sample

Table S5. Percent change in placental mtDNA content in association with prenatal NO₂ exposure in INMA mothers that spent ≥ 15 hours a day at home

Table S6. Association between prenatal NO₂ and birth weight in INMA mothers that spent ≥ 15 hours a day at home